



Burkhart Pediatric & Adolescent Dermatology

(919) 476-1118 | (855) 878-6037

135 Parkway Office Court - Suite 201

Cary, NC 27518

info@burkhartpediatricdermatology.com

www.burkhartpediatricdermatology.com

What is an excision of the skin?

An excision is a minimally invasive method of sampling the skin in order to make a diagnosis. An excision removes all or part of a skin lesion.

What is done during the test?

During the test the area of skin where the excision will be done is identified and cleaned well. You will be given a local anesthetic with a shot near the area. You may feel a prick from the shot. Once the area is numb a pencil-eraser sized piece of skin is removed. The area may need to be closed with sutures. The sutures will need to be removed in 5-7 (on the face and groin) or 10-14 days (on the scalp or body).

How do you care for the site after the excision?

Gently remove your dressing the morning after your procedure. Change the dressing at least once a day and if it becomes wet.

Please follow these guidelines for daily wound

care:

- Wash your hands with soap and water.
- Gently clean the area with soap and water using a cotton-tipped swab or gauze. (If you cannot easily remove the crust, soak the area with a wet piece of gauze for 15-20 minutes).
- After removing crusty material, gently pat the area dry with clean gauze.
- Use a clean cotton-tipped swab to apply a thick layer of petroleum jelly.
- Cover the area with clean gauze and tape or a bandage.
- Continue this process daily until you have your sutures removed.

Additional Instructions

- Avoid soaking in water until 14 days after your excision. You may shower as normal the morning after the excision.
- Pain at the excision site is typically minimal, but you may take over-the-counter pain relievers if needed.
- A minor amount of blood from the excision site is normal for a few days after the excision. If your wound bleeds enough to soak the gauze follow these instructions:
 - Place clean gauze on top of the bloody bandage and do not remove the bandage.
 - Apply firm, continuous pressure with the gauze for 20 minutes. If bleeding continues, apply pressure for another 20 minutes. Firm pressure should be slightly painful.
 - If bleeding stops wait a few hours then gently replace the bandage with a clean one.
 - If the bleeding continues after 40 minutes of pressure, contact us right away at (919) 476-1118.

Watch for Signs of Infection

Infections are very rare with excisions. If you develop the following symptoms or feel that your wound may be infected please contact us at (919) 476-1118:

- Increasing pain, soreness, tenderness, redness, or warmth around the area after the second day; worsening pain is frequently the first sign,
- A temperature of 100.4 degrees or greater,
- Chills
- Drainage from the wound that looks white, yellow-green or foul smelling.

Suture Removal

Depending on the type of suture used on your wound you may need to come back for suture removal in 5-7 days (on the face and groin) or 10-14 days (on the scalp or body). If you would prefer to have your sutures removed by one of our team members, please schedule the appointment the same day as the procedure. You may also have the sutures removed by your primary care doctor.